

A GAUSS-KUZMIN THEOREM FOR SOME CONTINUED FRACTION EXPANSIONS

Dan Lascu*

Mircea cel Batran Naval Academy, 1 Fulgerului, 900218 Constanta, Romania

Katsunori Kawamura†

College of Science and Engineering, Ritsumeikan University,

1-1-1 Noji Higashi, Kusatsu, Shiga 525-8577, Japan

August 24, 2011

Abstract

We consider a family of continued fraction expansions of any number in the unit closed interval $[0, 1]$ whose digits are differences of consecutive non-positive integer powers of an integer $m \geq 2$. For this expansion, we apply the method of Rockett and Szűsz from [6] and obtained the solution of its Gauss-Kuzmin type problem.

Mathematics Subject Classifications (2010). 11J70, 11K50

Key words: continued fractions, invariant measure, Gauss-Kuzmin problem

1 Introduction

The purpose of this paper is to prove a Gauss-Kuzmin type problem for non-regular continued fraction expansions introduced by Chan [1].

1.1 Gauss' Problem

One of the first and still one of the most important results in the metrical theory of continued fractions is so-called Gauss-Kuzmin theorem. Write $x \in [0, 1]$ as a regular continued fraction

$$x = \frac{1}{a_1 + \frac{1}{a_2 + \frac{1}{a_3 + \ddots}}} := [a_1, a_2, a_3, \dots],$$

*e-mail: lascudan@gmail.com.

†e-mail: kawamura@kurims.kyoto-u.ac.jp.

where $a_n \in \mathbb{N}_+ := \{1, 2, 3, \dots\}$. The metrical theory of continued fractions started on 25th October 1800, with a note by Gauss in his mathematical diary. Gauss wrote that (in modern notation)

$$\lim_{n \rightarrow \infty} \lambda(\tau^n \leq x) = \frac{\log(1+x)}{\log 2}, \quad x \in I := [0, 1].$$

Here λ is Lebesgue measure and the map $\tau : [0, 1) \rightarrow [0, 1)$, the so-called *regular continued fraction* (or *Gauss*) transformation, is defined by

$$\tau(x) := \frac{1}{x} - \left\lfloor \frac{1}{x} \right\rfloor, \quad x \neq 0; \quad \tau(0) := 0,$$

where $\lfloor \cdot \rfloor$ denotes the *floor* (or *entire*) function. Gauss' proof (if any) has never been found. A little more than 11 years later, in a letter dated 30 January 1812, Gauss asked Laplace to estimate the error

$$e_n(x) := \lambda(\tau^{-n}[0, x]) - \frac{\log(1+x)}{\log 2}, \quad n \geq 1, \quad x \in I.$$

This has been called *Gauss' Problem*. It received a first solution more than a century later, when R.O. Kuzmin (see [3]) showed in 1928 that $e_n(x) = \mathcal{O}(q^{\sqrt{n}})$ as $n \rightarrow \infty$, uniformly in x with some (unspecified) $0 < q < 1$. One year later, using a different method, Paul Lévy (see [5]) improved Kuzmin's result by showing that $|e_n(x)| \leq q^n$, $n \in \mathbb{N}_+$, $x \in I$, with $q = 3.5 - 2\sqrt{2} = 0.67157\dots$. The Gauss-Kuzmin-Lévy theorem is the first basic result in the rich metrical theory of continued fractions.

1.2 A non-regular continued fraction expansion

In this paper, we consider a generalization of the Gauss transformation and prove an analogous result. This transformation was studied in detail by Chan in [1] and Lascu and Kawamura in [4].

In [1], Chan shows that any $x \in [0, 1)$ can be written in the form

$$x = \frac{m^{-a_1(x)}}{1 + \frac{(m-1)m^{-a_2(x)}}{1 + \frac{(m-1)m^{-a_3(x)}}{1 + \ddots}}} := [a_1(x), a_2(x), a_3(x), \dots]_m, \quad (1.1)$$

where $m \in \mathbb{N}_+$, $m \geq 2$ and $a_n(x)$'s are non-negative integers.

For any $m \in \mathbb{N}_+$ with $m \geq 2$, define the transformation τ_m on I by

$$\tau_m(x) = \begin{cases} \frac{m^{\left\{\frac{\log x^{-1}}{\log m}\right\}} - 1}{m-1}, & \text{if } x \neq 0 \\ 0, & \text{if } x = 0, \end{cases} \quad (1.2)$$

where $\{\cdot\}$ stands for fractionary part. It is easy to see that τ_m maps the set Ω of irrationals in I into itself. For any $x \in (0, 1)$ put

$$a_n(x) = a_1(\tau_m^{n-1}(x)), \quad n \in \mathbb{N}_+,$$

with $\tau_m^0(x) = x$ and

$$a_1(x) = \begin{cases} \lfloor \log x^{-1} / \log m \rfloor, & \text{if } x \neq 0 \\ \infty, & \text{if } x = 0. \end{cases}$$

Transformation τ_m which generates the continued fraction expansion (1.1) is ergodic with respect to an invariant probability measure, γ_m , where

$$\gamma_m(A) = k_m \int_A \frac{dx}{((m-1)x+1)((m-1)x+m)}, \quad A \in \mathcal{B}_I, \quad (1.3)$$

with $k_m = \frac{(m-1)^2}{\log(m^2/(2m-1))}$ and $\mathcal{B}_I$ is the σ -algebra of Borel subsets of I (which, by definition, is the smallest σ -algebra containing intervals).

Next, we briefly present the results of Lascu and Kawamura in [4]. Were given some basic metric properties of the continued fraction expansion in (1.1). Hence, it was found the Brodén-Borel-Lévy formula used to determine the probability structure of $(a_n)_{n \in \mathbb{N}_+}$ under λ (=the Lebesgue measure).

Also, it was given the so-called *natural extension* $\bar{\tau}_m$, and were defined and studied the extended incomplete quotients $\bar{a}_l$, $l \in \mathbb{Z}$.

The associated Perron-Frobenius operator under different probability measures on $\mathcal{B}_I$ was derived. The Perron-Frobenius operator of τ_m under the invariant measure γ_m induced by the limit distribution function was studied and it was derived the asymptotic behaviour of this operator. Also, the Perron-Frobenius operator was restricted to the linear space of all complex-valued functions of bounded variation and to the space of all bounded measurable complex-valued functions.

The main result in [4] is the solving of Gauss-Kuzmin type problem using the method of random systems with complete connections by Iosifescu [2].

1.3 Main theorem

We show our main theorem in this subsection. For this purpose let μ be a non-atomic probability measure on $\mathcal{B}_I$ and define

$$\begin{aligned} F_n(x) &= \mu(\tau_m^n < x), \quad x \in I, \quad n \in \mathbb{N}, \\ F(x) &= \lim_{n \rightarrow \infty} F_n(x), \quad x \in I, \end{aligned} \quad (1.4)$$

with $F_0(x) = \mu([0, x))$. Our main result is the following theorem.

Theorem 1.1. *Let $k_m = \frac{(m-1)^2}{\log(m^2/(2m-1))}$. Then there exists*

$$F_n(x) = \frac{k_m}{(m-1)^2} \log \frac{m((m-1)x+1)}{(m-1)x+m} + \mathcal{O}(q_m^n), \quad (1.5)$$

where $0 < q_m < 1$.

2 Proof of Theorem 1.1

First, we show that $\{F_n\}$, defined in (1.4), satisfy a Gauss-Kuzmin-type equation, i.e., the following holds.

Proposition 2.1. *If $\{F_n\}$ are the functions defined in (1.4), then F_n satisfies the following Gauss-Kuzmin-type equation*

$$F_{n+1}(x) = \sum_{i \in \mathbb{N}} \left\{ F_n(\alpha^i) - F_n\left(\frac{\alpha^i}{1 + (m-1)x}\right) \right\}, \quad x \in I, \quad n \in \mathbb{N}. \quad (2.1)$$

Proof. Let $\alpha = 1/m$. Since $\tau_m^n(x) = \frac{m^{-a_{n+1}(x)}}{1 + (m-1)\tau_m^{n+1}(x)}$ it follows that

$$\begin{aligned} F_{n+1}(x) &= \mu(\tau_m^{n+1} < x) = \sum_{i \in \mathbb{N}} \mu\left(\frac{\alpha^i}{1 + (m-1)x} < \tau_m^n < \alpha^i\right) \\ &= \sum_{i \in \mathbb{N}} \left(F_n(\alpha^i) - F_n\left(\frac{\alpha^i}{1 + (m-1)x}\right) \right). \end{aligned}$$

□

Assuming that for some $p \in \mathbb{N}$ the derivative F'_p exists everywhere in I and is bounded, it is easy to see by induction that F'_{p+n} exists and is bounded for all $n \in \mathbb{N}_+$. This allows us to differentiate (1.4) term by term, obtaining

$$F'_{n+1}(x) = \sum_{i \in \mathbb{N}} \frac{(m-1)\alpha^i}{(1 + (m-1)x)^2} F'_n\left(\frac{\alpha^i}{1 + (m-1)x}\right). \quad (2.2)$$

Further, write $f_n(x) = (1 + (m-1)x)(m + (m-1)x)F'_n(x)$, $x \in I$, $n \in \mathbb{N}$, then (2.2) is rewritten as

$$f_{n+1}(x) = \sum_{i \in \mathbb{N}} P_m^i((m-1)x) f_n\left(\frac{\alpha^i}{1 + (m-1)x}\right), \quad (2.3)$$

where

$$P_m^i(x) = \frac{(m-1)\alpha^{i+1}(x+1)(x+m)}{(x + (m-1)\alpha^i + 1)(x + (m-1)\alpha^{i+1} + 1)}, \quad x \in I. \quad (2.4)$$

Putting $\Delta_i = \alpha^i - \alpha^{2i}$, $i \in \mathbb{N}$, we get

$$\begin{aligned} P_m^i((m-1)x) &= (m-1) \left[\alpha^{i+1} + \frac{\Delta_i}{(m-1)x + (m-1)\alpha^i + 1} \right. \\ &\quad \left. - \frac{\Delta_{i+1}}{(m-1)x + (m-1)\alpha^{i+1} + 1} \right]. \end{aligned}$$

Proof of Theorem 1.1 Introduce a function $R_n(x)$ such that

$$F_n(x) = \frac{k_m}{(m-1)^2} \log \frac{m((m-1)x+1)}{(m-1)x+m} + R_n\left(\frac{k_m}{(m-1)^2} \log \frac{m((m-1)x+1)}{(m-1)x+m}\right), \quad (2.5)$$

where $k_m = \frac{(m-1)^2}{\log(m^2/(2m-1))}$.

Because $F_n(0) = 0$ and $F_n(1) = 1$, we have $R_n(0) = R_n(1) = 0$. To prove the theorem, we have to show that

$$R_n(x) = \mathcal{O}(q_m^n), \quad (2.6)$$

where $0 < q_m < 1$.

If we can show that $f_n(x) = k_m + \mathcal{O}(q_m^n)$, then its integration will show the equation (1.5).

To demonstrate that $f_n(x)$ has this desired form, it suffices to establish that $f'_n(x) = \mathcal{O}(q_m^n)$.

We have

$$\begin{aligned} (P_m^i((m-1)x))' &= (m-1) \left[\frac{(m-1)\Delta_{i+1}}{((m-1)x + (m-1)\alpha^{i+1} + 1)^2} \right. \\ &\quad \left. - \frac{(m-1)\Delta_i}{((m-1)x + (m-1)\alpha^i + 1)^2} \right]. \end{aligned}$$

Now from (2.3), we have

$$\begin{aligned} f'_{n+1}(x) &= \sum_{i \in \mathbb{N}} \left[(P_m^i((m-1)x))' f_n \left(\frac{\alpha^i}{1 + (m-1)x} \right) \right. \\ &\quad \left. - P_m^i((m-1)x) \frac{(m-1)\alpha^i}{((m-1)x + 1)^2} f'_n \left(\frac{\alpha^i}{1 + (m-1)x} \right) \right] \\ &= -(m-1) \sum_{i \in \mathbb{N}} (A_i + B_i) \end{aligned} \quad (2.7)$$

where

$$\begin{aligned} A_i &= \frac{(m-1)\Delta_{i+1}}{((m-1)x + (m-1)\alpha^{i+1} + 1)^2} \\ &\quad \times \left(f_n \left(\frac{\alpha^{i+1}}{(m-1)x + 1} \right) - f_n \left(\frac{\alpha^i}{(m-1)x + 1} \right) \right) \\ B_i &= P_m^i((m-1)x) \frac{(m-1)\alpha^i}{((m-1)x + 1)^2} f'_n \left(\frac{\alpha^i}{1 + (m-1)x} \right). \end{aligned}$$

By applying the mean value theorem of calculus to the difference

$$f_n \left(\frac{\alpha^{i+1}}{(m-1)x + 1} \right) - f_n \left(\frac{\alpha^i}{(m-1)x + 1} \right),$$

we obtain

$$f_n \left(\frac{\alpha^{i+1}}{(m-1)x + 1} \right) - f_n \left(\frac{\alpha^i}{(m-1)x + 1} \right) = \left\{ \frac{\alpha^{i+1}}{(m-1)x + 1} - \frac{\alpha^i}{(m-1)x + 1} \right\} f'_n(\theta_i),$$

with

$$\frac{\alpha^{i+1}}{(m-1)x+1} < \theta_i < \frac{\alpha^i}{(m-1)x+1}.$$

Thus, from (2.7), we have

$$\begin{aligned} f'_{n+1}(x) &= (m-1)^2 \sum_{i \in \mathbb{N}} \frac{\alpha^{i+1} \Delta_{i+1}}{((m-1)x+1)((m-1)x+(m-1)\alpha^{i+1}+1)^2} f'_n(\theta_i) \\ &\quad - (m-1) \sum_{i \in \mathbb{N}} P_m^i((m-1)x) \frac{\alpha^i}{((m-1)x+1)^2} f'_n\left(\frac{\alpha^i}{1+(m-1)x}\right). \end{aligned} \quad (2.8)$$

Let M_n be the maximum of $|f'_n(x)|$ on I , i.e., $M_n = \max_{x \in I} |f'_n(x)|$. Then (2.8) implies

$$\begin{aligned} M_{n+1} &\leq M_n \cdot \max_{x \in I} \left| (m-1)^2 \sum_{i \in \mathbb{N}} \frac{\alpha^{i+1} \Delta_{i+1}}{((m-1)x+1)((m-1)x+(m-1)\alpha^{i+1}+1)^2} \right. \\ &\quad \left. + (m-1) \sum_{i \in \mathbb{N}} P_m^i((m-1)x) \frac{\alpha^i}{((m-1)x+1)^2} \right|. \end{aligned} \quad (2.9)$$

We now must calculate the maximum value of the sums in this expression.

First, we note that

$$\frac{\alpha^{i+1} \Delta_{i+1}}{((m-1)x+1)((m-1)x+(m-1)\alpha^{i+1}+1)^2} \leq \frac{\alpha^{2i+2}}{((m-1)\alpha^{i+1}+1)^2}. \quad (2.10)$$

Note that, in the upper inequality, we have used that $\Delta_i = \alpha^i - \alpha^{2i}$ and that $0 \leq x \leq 1$.

Next, observe that the function

$$h_m(x) := \frac{\alpha^i P_m^i((m-1)x)}{((m-1)x+1)^2}$$

is decreasing for $x \in I$ and $i \in \mathbb{N}$. Hence, $h_m(x) \leq h_m(0)$. This leads to

$$\frac{\alpha^i P_m^i((m-1)x)}{((m-1)x+1)^2} \leq \frac{(m-1)\alpha^{2i}}{((m-1)\alpha^{i+1}+1)^2}. \quad (2.11)$$

The relations (2.10) and (2.11) allow us to rewrite inequality (2.9) as

$$M_{n+1} \leq q_m \cdot M_n, \quad (2.12)$$

where

$$q_m := (m-1)^2(m^2+1) \sum_{i \in \mathbb{N}} \frac{1}{(m^{i+1}+m-1)^2}. \quad (2.13)$$

Since, for $i \geq 2$, we have

$$\frac{1}{(m^{i+1}+m-1)^2} \leq \frac{1}{m^2(m-1)^2(m^2+1)} \left(\frac{1}{m}\right)^i,$$

therefore,

$$\begin{aligned}
q_m &\leq (m-1)^2(m^2+1) \\
&\quad \times \left(\frac{1}{(2m-1)^2} + \frac{1}{m^2+m-1} + \frac{1}{m^2(m-1)^2(m^2+1)} \sum_{i \geq 2} \left(\frac{1}{m} \right)^i \right) \\
&= (m-1)^2(m^2+1) \left(\frac{1}{(2m-1)^2} + \frac{1}{m^2+m-1} + \frac{1}{m^3(m-1)^3(m^2+1)} \right) \\
&\leq 1,
\end{aligned}$$

for any $m \in \mathbb{N}$, $m \geq 2$. □

References

- [1] Chan, H.-C., *The asymptotic growth rate of random Fibonacci type sequences. II*, Fibonacci Quart.**44** (2006), 73-84.
- [2] Iosifescu, M. and Grigorescu, S., *Dependence With Complete Connections and its Applications*, Cambridge Tracts in Mathematics **96**, 1990. Cambridge Univ.Press, Cambridge. [(2009): second printing slightly corrected].
- [3] Kuzmin, R.O., *On a problem of Gauss*, Dokl. Akad. Nauk SSSR Ser. A (1928) 375-380. [Russian; French version in *Atti Congr. Internaz.Mat. (Bologna, 1928)*, Tomo **VI** (1932) 83-89. Zanichelli, Bologna].
- [4] Lascu, D. and Katsunori, K., *On a Gauss-Kuzmin-type problem for a family of continued fraction expansions*, arXiv:1108.3441v1 (2011), 1-32.
- [5] Lévy, P., *Sur les lois de probabilité dont dépendent les quotients complets et incomplets d'une fraction continue*, Bull. Soc. Math. France **57** (1929), 178-194.
- [6] Rockett, A.M. and Szűsz, P., *Continued Fractions*, World Scientific, Singapore, 1992.